Appendix E: Habitat and Species Data informing the Risk Assessment

This Appendix is provided in support to the following report:

Marine Planning Consultants Ltd. (2014). Lyme Bay Fisheries and Conservation Reserve: Integrated Fisheries Management Plan. A report produced for the Lyme Bay Fisheries and Conservation Reserve Working Group, UK.

The report, submitted 18/09/2014, addresses comments made by the wider Lyme Bay Fisheries and Conservation Reserve Working Group at a Workshop 09/09/2014.

Contents

Source of data1
Species Inventory & Conservation Status
All species identified3
Species of conservation importance
Selected species of conservation importance
Figures
Figure E1: Location of benthic species selected for risk assessment (Latin names)
Tables
Table E1. Source of data for each feature type1
Table E2. Species data inventory: species of conservation importance4

Source of data

Table E1. Source of data for each feature type

Sector	Information Type	Source Project	Source Organisation	Data Type	Comment
BENTHIC HABITATS	Biotope model	Various	Devon Wildlife Trust (Devon Biodiversity Records Centre)	Full coverage interpolation	Map provided in report
BENTHIC HABITATS	Reef map	cSAC characterisation	Natural England / Cefas	GIS model	Map provided in report
BENTHIC SPECIES	Diving observations, benthic grabs and video transects	Various	Devon Wildlife Trust (Devon Biodiversity Records Centre)	Point data	Map and table of species of conservation importance provided in report
BENTHIC SPECIES	Various	Various	NBN	Point data	Map and table of species of conservation importance provided in report
BENTHIC SPECIES	Diving observations	Various	Seasearch	Point data	Map and table of species of conservation importance provided in report
BENTHIC SPECIES	Video transects	Impacts of scallop dredging on epibenthic fauna	Bangor University / NERC	Point data	Table of species of conservation importance provided in report
FISH	Fish Distribution	Nursery and spawning areas	Cefas (Ellis, et al., 2012)	Gridded	Map and table of species of conservation importance provided in report
FISH	Fish Distribution	Various	Devon Wildlife Trust (Devon Biodiversity Records Centre)	Point data	Table of species of conservation importance provided in report
FISH	Fish Distribution	N/A	MARINE-Life	Point data	Not provided in time to be included in report (but will be assessed)
FISH	Fish Distribution	NBN	JNCC	Point data	Table of species of conservation importance provided in report
FISH	Fish Distribution	Various	Devon Wildlife Trust (Devon Biodiversity Records Centre)	Point data	Table of species of conservation importance provided in report

Sector	Information Type	Source Project	Source Organisation	Data Type	Comment
CETACEAN	Cetacean sightings	N/A	MARINE-Life	Point data	Not provided in time to be included in report (but will be assessed)
CETACEAN	Cetacean sightings	SCANS-II	University of St Andrews	Point data	Not in AOI
TURTLES	Turtle sightings	Various	Devon Wildlife Trust (Devon Biodiversity Records Centre)	Point data	Table of species of conservation importance provided in report
TURTLES	Turtle sightings	N/A	MARINE-Life	Point data	Not provided in time to be included in report (but will be assessed)
TURTLES	Turtle sightings	N/A	NBN Gateway	Point data	Map and table of species of conservation importance provided in report
BIRDS	Bird sightings	N/A	MARINE-Life	Point data	Not provided in time to be included in report (but will be assessed)
BIRDS	Bird breeding, wintering and moulting sightings	N/A	JNCC	Point data	Map and table of species of conservation importance provided in report
BIRDS	Bird sightings	European Seabirds at Sea (ESAS) and Seabird Monitoring Programme (SMP)	JNCC	Point data	Map and table of species of conservation importance provided in report

Species Inventory & Conservation Status

All species identified

A species inventory that lists all species identified in the Lyme Bay AOI is available in Excel spreadsheet format: please request the Lyme Bay species inventory spreadsheet. This table details conservation status; species and habitats record data source; characteristics, e.g. habitat and feeding; and detail of records for spatial evidence in Lyme Bay, e.g. vintage, rocky reef association. Please request the Lyme Bay conservation tables.

Species of conservation importance

The summary table below (Table E1) shows all species identified in the Lyme Bay AOI that have a conservation status, with each type shown. Justification is given for those that were selected for the risk assessment.

Table 1E shows a dark blue highlight where the species / habitat was selected for assessment in the project and assigned 'Tier 1'. Light blue highlights also indicate species / habitat selected although these are have only a few records and / or are old records. Due to their significant conservation status, e.g. to protect against rarity, they have been included in this Appendix and are discussed in the main report; however these are not of priority in the assessment and are therefore assigned 'Tier 2', and may be used to help inform future monitoring.

Note the Latin name is the WoRMS¹ software accepted name; and species names are not in Italics due to the original spreadsheet format.

¹ http://www.marinespecies.org/

 Table E2. Species data inventory: species of conservation importance

Species Nam	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Benthic / Epibenthic	Sabellaria alveolata	Honeycomb Worm	Habitats Directive, English NERC List	The majority of observations are recent (last 20 years). There are a high number of observations of the species (46) most of which report high abundances which could be indicative of reef (S, A, or C) - also observations are likely to have been of crust or reef since anything smaller is less likely to be encountered. Although S. alveolata observations were exclusively associated with the rocky reef there is potential for it to occur outside of the protected rocky reef area. Whilst it is not certain whether the records indicate reef, if they do then the habitat is protected in its own right it should be treated as a separate habitat. However due to the lack of evidence as to whether it is or is not reef forming in Lyme Bay, and that it is in an area that is widely surveyed due to conservation interest yet has still not been officially observed, this can only be included as a Tier 2 habitat - as a precautionary approach.	SABELLARIA	Tier 2
Benthic / Epibenthic	Sabellaria spinulosa	Ross Worm	Habitats Directive, OSPAR (2008), English NERC List	The majority of observations are recent (last 20 years) and there are a high number of observations (32). Although only a small number of the observations were of high abundances this may in part be a reflection of fishing pressures already being exerted on this area and there may therefore be potential for more reef habitats to develop in the future. Although many of the S. spinulosa observations were associated with the rocky reef there is potential for this habitat to occur outside of the protected rocky reef area. Whilst it is not certain whether the records indicate reef, if they do then the habitat is protected in its own right it should be treated as a separate habitat. However due to the lack of evidence as to whether it is or is not reef forming in Lyme Bay, and that it is in an area that is widely surveyed due to conservation interest yet has still not been officially observed, this can only be included as a Tier 2 habitat - as a precautionary approach. Note that Sabellaria sinulosa is less common in reef form than Sabellaria alveolata.	SABELLARIA	Tier 2
Benthic / Epibenthic	Schizobrachiella sanguinea	A bryozoan	UK Cons. Status (Rare & Scarce)	A small number of recent records were associated exclusively with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Pycnoclavella aurilucens	Orange Lights Seasquirt	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Phallusia mammillata	A sea squirt	UK Cons. Status (Rare & Scarce)	A very high number (>400) of recent (last 20 years) high abundance records (S,A or C) the majority of which were associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 1
Benthic / Epibenthic	Aiptasia mutabilis	Trumpet Anemone	UK Cons. Status (Rare & Scarce)	A very high number (>200) of recent (last 20 years) high abundance records (S,A or C) the majority of which were associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 1

Species Nam	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Benthic / Epibenthic	Amphianthus dohrnii	Sea-fan Anemone	English NERC List, UK Cons. Status (Rare & Scarce)	Only a single record of this very tiny species (which is likely to be overlooked by most sampling techniques). Species is exclusively associated with the protected rocky reef habitat and more specifically with the sea fans and will therefore be assessed under the umbrella habitat of rocky reef and does not requires additional consideration.	REEF & PINK SEAFAN	Tier 2
Benthic / Epibenthic	Eunicella verrucosa	Pink Sea Fan	OSPAR (2008), IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, UK Cons. Status (Rare & Scarce)	There are a very large number of records of this species (254) although this is likely to be inpart due to sampling biases, since a number of the surveys have been specifically targeting this species, and also its conspicuous nature. The majority of records are recent and indicate that this species is strongly associated with the protected rocky reef. However a relatively large number (53) of observations of this species in non-reef areas indicates that there is a significant seafan population associated with other habitats within the AOI. Given the high conservation status of this species (listed as vulnerable on the IUCN Red List as well as T&D Ospar list) the sensitivity of this species will be assessed in its own right as well as under the umbrella of the rocky reef.	REEF & PINK SEAFAN	Tier 1
Benthic / Epibenthic	Caryophyllia (Caryophyllia) inornata	Southern Cup Coral	OSPAR (2008), UK Cons. Status (Rare & Scarce)	A high number (32) of recent observations strongly associated with the protected rocky reefs some of which indicate that this species is common or frequent. Although there are records of this species outside the protected rocky reef feature, this species is only found on rock which is not mapped elsewhere in the AOI. Therefore these records are likely to indicate the presence of very small rocky features and/or areas of rocky reef that have not yet been mapped. This species will therefore be assessed under the umbrella habitat of rocky reefs and the locations of observations outwith the reef will be provided to the IFCAs to inform future survey efforts.	REEF	Tier 1
Benthic / Epibenthic	Caryophyllia (Caryophyllia) smithii	Devonshire Cup Coral	OSPAR (2008)	A very high number (233) of recent observations strongly associated with the protected rocky reefs most of which indicate that this species is occasional or rare. Although there are records of this species outside the protected rocky reef feature, this species is predominantly only found on rock which is not mapped elsewhere in the AOI. Therefore these records are likely to indicate the presence of very small rocky features and/or areas of rocky reef that have not yet been mapped. This species will therefore be assessed under the umbrella habitat of rocky reefs and the locations of observations outwith the reef will be provided to the IFCAs to inform future survey efforts.	REEF	Tier 1
Benthic / Epibenthic	Hoplangia durotrix	Weymouth carpet Coral	OSPAR (2008), UK Cons. Status (Rare & Scarce)	A small number (7) of recent observations strongly associated with the protected rocky reefs most of which indicate that this species is rare. Although there are records of this species outside the protected rocky reef feature, this species is only found on rock which is not mapped elsewhere in the AOI. Therefore these records are likely to indicate the presence of very small rocky features and/or areas of rocky reef that have not yet been mapped. This species will therefore be assessed under the umbrella habitat of rocky reefs and the locations of observations outwith the reef will be provided to the IFCAs to inform future survey efforts.	REEF	Tier 2

Species Nam	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Benthic / Epibenthic	Leptopsammia pruvoti	Sunset Cup Coral	OSPAR (2008), English NERC List, UK Cons. Status (Rare & Scarce)	A moderate number (22) of recent observations strongly associated with the protected rocky reefs most of which indicate that this species is abundant. Although there are records of this species outside the protected rocky reef feature, this species is only found on rock which is not mapped elsewhere in the AOI. Therefore these records are likely to indicate the presence of very small rocky features and/or areas of rocky reef that have not yet been mapped. This species will therefore be assessed under the umbrella habitat of rocky reefs and the locations of observations outwith the reef will be provided to the IFCAs to inform future survey efforts.	REEF	Tier 1
Benthic / Epibenthic	Aglaophenia kirchenpaueri	A hydroid	UK Cons. Status (Rare & Scarce)	A single recent observation has been made of this species within the AOI. Although this record does not fall within the protected rocky reef area it is closely adjacent (50m). As this observation was made by a diver the positional accuracy of this observation is cannot be assured. This species occurs on rock and therefore it is thought highly probable that this species is associated with the rocky reef and it will be assessed as such. The position will also be provided to the IFCAs as a focus for future survey work to further refine the reef boundaries and / or identify any other areas of rocky habitat.	REEF	Tier 2
Benthic / Epibenthic	Echinus esculentus	Edible Sea Urchin	IUCN Red List	A small number of recent records indicate that this species is exclusively associated with the protected rocky reef habitats. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Paracentrotus lividus	Purple Sea Urchin	UK Cons. Status (Rare & Scarce)	A small number of recent records indicate that this species is exclusively associated with the protected rocky reef habitats. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Ophiopsila aranea	A brittlestar	UK Cons. Status (Rare & Scarce)	A small number of recent records indicate that this species is exclusively associated with the protected rocky reef habitats. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Pholadidea loscombiana	Paper Piddock	UK Cons. Status (Rare & Scarce)	A small number of recent records indicate that this species is strongly associated with the protected rocky reef habitats. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration.	REEF	Tier 2
Benthic / Epibenthic	Modiolus modiolus	Horse Mussel	Habitats Directive, English NERC List	Very few records with low abundance, all of individual mussels. No evidence of beds that would qualify for protection and these are very unlikely to occur this far south. Therefore this species does not require any further consideration in the sensitivity modelling.	Excluded	N/A
Benthic / Epibenthic	Mytilus edulis	Blue Mussel	Habitats Directive, OSPAR (2008), English NERC List	A very high number of records (104) some of which may be indicative of the presence of mussel beds (A/C/F) which could qualify for protection. Mytilus beds are known to occur in the wider Lyme Bay. The majority of the records (~90%) are old (>20years) although the most recent record was made in 2004. The spatial representation of this habitat feature is not likely to be overly reliable and the locations of observations (particularly those that are recorded as A/C/F) will be supplied to the IFCAs as potential targets for future survey work. As there is a lack of evidence as to whether it is or is not forming mussel beds in Lyme Bay, and that it is in an area that is widely surveyed due to conservation interest yet has still not been officially observed, this can only be included as a Tier 2 habitat - as a precautionary approach - to reflect its conservation status and association not only with the rocky reef but also beyond this delineated area (25%).	MYTILUS	Tier 2

Species Nam	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Benthic / Epibenthic	Ostrea edulis	Native oyster	OSPAR (2008), English NERC List	Although only a small number of observations were made of this species (6) over five separate years up to 2004, it is relatively cryptic and could easily be overlooked by video and diver surveys. There is a partial association with the protected rocky reefs but given the high conservation status of this species (as both a species and as a habitat where it forms beds) it will be assessed in its own right, though as Tier 2 assessment.	NATIVE OYSTER	Tier 2
Benthic / Epibenthic	Arctica islandica	Icelandic Cyprine / Ocean Quahog	OSPAR (2008)	Although only a small number of observations were made of this species (4) it is could easily be overlooked by video and diver surveys as it is usually buried or partially buried. Given the high conservation status of this species and its longevity, it will be assessed in its own right.	OCEAN QUAHOG	Tier 2
Benthic / Epibenthic	Nucella lapillus	Dog Whelk	OSPAR (2008)	A high number of records, some of which are recent, were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 1
Benthic / Epibenthic	Trapania pallida	A sea slug	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 2
Benthic / Epibenthic	Tritonia nilsodhneri	A sea slug	UK Cons. Status (Rare & Scarce)	A large number of recent records were strongly associated with the protected rocky reef. This species is commensurate with the Pink Seafan and will therefore be assessed under the umbrella Pink Seafan and does not require additional consideration.	PINK SEAFAN	Tier 1
Benthic / Epibenthic	Dysidea pallescens	A sponge	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 2
Benthic / Epibenthic	Adreus fascicularis	Branched yellow sponge	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 2
Benthic / Epibenthic	Axinella damicornis	A sponge	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 2
Benthic / Epibenthic	Tethyspira spinosa	Pork scratching sponge	UK Cons. Status (Rare & Scarce)	A small number of recent records were strongly associated with the protected rocky reef. This species will therefore be assessed under the umbrella habitat of the rocky reef and does not require additional consideration. Geographic positions of records from outside the reef will be supplied to the IFCAs as potential targets for future survey efforts.	REEF	Tier 2

Species Na	me / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Algae & Ichen	Zanardinia typus	Penny Weed	UK Cons. Status (Rare & Scarce)	A small number of recent records, some of which are associated with the protected rocky reef. Although there are some records outwith the rocky reef, this species is usually found on rock and hence, the additional observations may represent isolated rock outcrops or areas of rocky reef that have not yet been accurately mapped. As these observations were made by divers it is also possible that there is some inaccuracy in the positioning. Therefore this species will be included in the sensitivity assessment under the umbrella of the rocky reef habitat and the coordinates of observations outwith the reef will be provided to the IFCAs as a potential target for future survey work.	REEF	Tier 2
Algae & Ichen	Padina pavonica	Peacocks Tail	English NERC List, UK Cons. Status (Rare & Scarce)	A small number of recent records all of which are associated with the protected rocky reef. Therefore this species will be included in the sensitivity assessment under the umbrella of the rocky reef habitat and no further consideration is required.	REEF	Tier 2
Algae & Ichen	Lithothamnion	Maerl	Habitats Directive, OSPAR (2008), English NERC List, UK Cons. Status (Rare & Scarce)	A small number of records indicate that Lithothamnion is abundant at some locations in the AOI indicating that beds may be present. Whilst calcareous algae have the capacity to form maerl deposits, the encrusting forms are more likely to found on boulders and stones, rather than the flattened leaf-like thalli that detach and accumulate to form a maerl habitat. Therefore presence of a potential maerl species such as Lithothamnion or Lithophyllum does not necessarily indicate areas of 'maerl'. However as this species is protected as both a habitat and a species it will be included in the sensitivity assessment as Maerl. All records are point records at present and hence the locations will be provided to the IFCAs as a potential target for future survey work.	MAERL	Tier 2
Algae & Ichen	Phymatolithon calcareum	Maerl	Habitats Directive, OSPAR (2008), English NERC List	A small number of records indicate that Phymatolithon calcareumis are abundant at some locations in the AOI indicating that beds may be present. As this species is protected as both a habitat and a species it will be included in the sensitivity assessment as Maerl. All records are point records at present and hence the locations will be provided to the IFCAs as a potential target for future survey work.	MAERL	Tier 2
Fish	Anguilla anguilla	European Eel	OSPAR (2008), EC-CITES , IUCN Red List, English NERC List	Only one record of the European Eel exists in the AOI and this record is 17 years old. The data available were not from targeted fisheries surveys and hence the abundance of fish species is likely to be dramatically underrepresented; however the fisheries catch data only report Conger Eels. Given the high conservation status and unusual life history of this species it will be assessed in the sensitivity modelling at the species level - though at Tier 2 due to the single / old record - but with the precautionary approach.	EUROPEAN EEL	Tier 2
Fish	Gadus morhua	Atlantic Cod	OSPAR (2008), IUCN Red List, English NERC List	Although there are only 15 records (1994 to 2004) it is likely that this species is dramatically underrepresented in the spatial data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7). This species will therefore be included at the species level.	COD	Tier 1
Fish	Merlangius merlangus	Whiting	English NERC List,	Although there are only a small number of records (1 from 2003) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7). This species will therefore be included at the species level.	WHITING	Tier 1

Species Na	me / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Fish	Molva molva	Ling	English NERC List	Although there are only a small number of records (4 from 1998-2009) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7); however this is very small, 25kg 2002-2013, albeit only recording individuals >63cm MLS so not juveniles etc. This species will therefore be included at the species level - though at Tier 2.	LING	Tier 2
Fish	Lophius piscatorius	Angler	English NERC List	Although there are only a small number of records (2 from 2007) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7). This species will therefore be included at the species level.	ANGLERFISH	Tier 2
Fish	Pomatoschistus minutus	Sand Goby	Bern Convention	Although this species is strongly associated with the protected rocky reef there is a high chance that it will occur elsewhere in the AOI and this association may in-part be due to sampling biases with many of the surveys being focussed on the reef habitats (20 of 22 records on reef, 1994 - 2004). Given the high conservation status of this species and its importance in marine food webs this species will be assessed in its own right. However, the fishery landings data does not confirm presence of the species within the AOI - and so it is included at Tier 2.	SAND GOBY	Tier 2
Fish	Pleuronectes platessa	Plaice	IUCN Red List, English NERC List	Although there are only a small number of records (3 from 1998-2002) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7). This species will therefore be included at the species level.	PLAICE	Tier 1
Fish	Solea solea	Sole	English NERC List	Although there are only a small number of records (9 from 1994-2004) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7). This species will therefore be included at the species level.	SOLE	Tier 1
Fish	Prionace glauca	Blue Shark	IUCN Red List, English NERC List	Highly transient species unlikely to be strongly associated with any one habitat within the AOI therefore this species will be assessed under the assessment unit of Sharks. However only one record (2012) so Tier 2.	SHARKS	Tier 2
Fish	Scyliorhinus stellaris	Nursehound	IUCN Red List	Highly transient species unlikely to be strongly associated with any one habitat within the AOI therefore this species will be assessed under the assessment unit of dogfish. 4 records 2006-2011, though not supported by fishery landings data.	DOGFISH	Tier 2
Fish	Alopias vulpinus	Thresher Shark	IUCN Red List	Highly transient species unlikely to be strongly associated with any one habitat within the AOI therefore this species will be assessed under the assessment unit of Sharks. However only one record (2010) so Tier 2.	SHARKS	Tier 2
Fish	Cetorhinus maximus	Basking shark	Bern Convention, Bonn Convention, OSPAR (2008), EC- CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981	Highly transient species unlikely to be strongly associated with any one habitat within the AOI . Very different life history to the other shark species recorded in this area therefore this species will be assessed at the species level. 8 records 1999-2011.	BASKING SHARK	Tier 1
Fish	Raja clavata	Thornback Ray	OSPAR (2008), IUCN Red List	Although there is only record (2003) it is likely that this species is dramatically underrepresented given the data types available. This species will therefore be included in the rays group given its high conservation status. However only one record so Tier 2.	RAYS	Tier 2

Species Nan	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Fish	Raja montagui	Spotted Ray	OSPAR (2008), IUCN Red List	Although there is only record (2012) it is likely that this species is dramatically underrepresented given the data types available. This species will therefore be included in the rays group given its high conservation status. However only one record so Tier 2.	RAYS	Tier 2
Fish	Squalus acanthias	Spurdog	Bonn Convention, OSPAR (2008), IUCN Red List, English NERC List	Although there is only record which is old (1977) it is likely that this species is dramatically underrepresented given the data types available. In support, the fishery landings data confirms presence of the species within the AOI (ICES rectangles 30E6 and 30E7) - 41kg 2002-2013. This species will therefore be included in the dogfish group given its high conservation status. However only one / old record and low landings weight so Tier 2.	DOGFISH	Tier 2
Mammals & Turtles	Balaenoptera acutorostrata	Minke whale	Bern Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although there is only record (2007), the distribution of marine mammals is under recorded throughout the UK. Whales are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Whales group.	WHALES	Tier 2
Mammals & Turtles	Balaenoptera physalus	Fin whale	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded once within the AOI (2000), the distribution of marine mammals is under recorded throughout the UK. Whales are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Whales group.	WHALES	Tier 2
Mammals & Turtles	Megaptera novaeangliae	Humpback whale	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES, IUCN Red List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded once within the AOI (1992), the distribution of marine mammals is under recorded throughout the UK. Whales are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Whales group.	WHALES	Tier 2
Mammals & Turtles	Globicephala melas	Long-finned pilot whale	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded twice within the AOI (2002, 2006), the distribution of marine mammals is under recorded throughout the UK. Whales are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Whales group.	WHALES	Tier 2
Mammals & Turtles	Grampus griseus	Risso's dolphin	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has not been recorded within the AOI, it has been recorded close by and the distribution of marine mammals is under recorded throughout the UK. Dolphins are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Dolphin and Porpoise group.	DOLPHINS & PORPOISES	Tier 2

Species Nan	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Mammals & Turtles	Delphinus delphis	Common dolphin	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI twelve times (1998-2012), the distribution of marine mammals is under recorded throughout the UK. Dolphins are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Dolphin and Porpoise group.	DOLPHINS & PORPOISES	Tier 1
Mammals & Turtles	Lagenorhynchus albirostris	White- beaked dolphin	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES, IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI once with three sightings (2008), the distribution of marine mammals is under recorded throughout the UK. Dolphins are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Dolphin and Porpoise group.	DOLPHINS & PORPOISES	Tier 2
Mammals & Turtles	Tursiops truncatus	Bottlenose dolphin	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Unlike the other dolphin and porpoise species there is some evidence to suggest that there is a resident population of Bottlenose dolphins within the AOI with 59 recorded sightings (1999-2012). This population of dolphins is likely to be more dependent on the AOI than other transient visitors and it is therefore included in the sensitivity modelling as an individual species. As there are 59 recorded sightings it may also be possible to undertake a spatial assessment for this species.	BOTTLENOSE DOLPHIN	Tier 1
Mammals & Turtles	Phocoena phocoena	Harbour porpoise	Bern Convention, Bonn Convention, Habitats Directive, OSPAR (2008), EC-CITES, IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI eighteen times (1998-2011), the distribution of marine mammals is under recorded throughout the UK. Dolphins are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Dolphin and Porpoise group.	DOLPHINS & PORPOISES	Tier 1
Mammals & Turtles	Halichoerus grypus	Grey seal	Bern Convention, Bonn Convention, Habitats Directive, IUCN Red List, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI four times (2008-2009), the distribution of marine mammals is under recorded throughout the UK. Seals are highly mobile and are likely to be transient visitors to this area, since there is no evidence of any haul out areas. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Seal group - Tier 1 due to the recent recordings, albeit low in number.	SEALS	Tier 1
Mammals & Turtles	Phoca vitulina	Harbour seal	Bern Convention, Bonn Convention, Habitats Directive, IUCN Red List, English NERC List, Cons. of Hab&Spp Regs 2010	Although this species has not been recorded within the AOI it has been recorded in adjacent areas and the distribution of marine mammals is under recorded throughout the UK. Seals are highly mobile and are likely to be transient visitors to this area, since there is no evidence of any haul out areas. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Seal group.	SEALS	Tier 2

Species Nan	ne / Group		Assessment			
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority
Mammals & Turtles	Caretta caretta	Loggerhead Turtle	Bern Convention, Bonn Convention, Habitats Directive, OSPAR (2008), EC-CITES, IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI once (1938) the distribution of turtles is under recorded throughout the UK. Turtles are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Turtle group.	TURTLES	Tier 2
Mammals & Turtles	Chelonia mydas	Green Turtle	Bern Convention, Bonn Convention, Habitats Directive, EC-CITES , IUCN Red List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI once and this is extremely old (1887) the distribution of turtles is under recorded throughout the UK. Turtles are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Turtle group.	TURTLES	Tier 2
Mammals & Turtles	Dermochelys coriacea	Leatherback turtle	Bern Convention, Bonn Convention, Habitats Directive, OSPAR (2008), EC-CITES, IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, Cons. of Hab&Spp Regs 2010	Although this species has only been recorded within the AOI once (2009) the distribution of turtles is under recorded throughout the UK. Turtles are highly mobile and are likely to be transient visitors to this area. However, they are subject to high levels of protection and do have the potential to interact with the inshore fisheries operating in this area. This species will therefore be included in the sensitivity modelling within the Turtle group.	TURTLES	Tier 2
Birds	Fulmarus glacialis	Northern Fulmar	Bern Convention, IUCN Red List, UK Cons. Status (Birds)	High number (74) or recent records and evidence that this species uses the area to breed therefore this species will be assessed in its own right.	FULMAR & SURFACE FEEDING BIRDS	Tier 1
Birds	Calonectris diomedea	Cory's Shearwater	Bern Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Very few records of this species (1) in the AOI which visits the area in the summer. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2
Birds	Hydrobates pelagicus	European Storm Petrel	Bern Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Very few records of this species (1) in the AOI which migrates through the area in spring and autumn. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2
Birds	Puffinus griseus	Sooty Shearwater	Bern Convention, IUCN Red List, UK Cons. Status (Birds)	Very few records of this species (2) in the AOI which migrates through the area in spring and autumn. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2
Birds	Puffinus mauretanicus	Balearic Shearwater	Bern Convention, Birds Directive, OSPAR (2008), IUCN Red List, English NERC List, UK Cons. Status (Birds)	Although there are only a small number of this species within the AOI there are reports that suggest this are may become increasingly important to this species and given its Red status it will be included in the sensitivity assessment at the species level.	BALEARIC SHEERWATRE & PURSUIT & PLUNGE DIVING BIRDS	Tier 1
Birds	Puffinus puffinus	Manx Shearwater	Bern Convention, IUCN Red List, UK Cons. Status (Birds)	Although there are a moderate number of records of the manx sheerwater in the AOI there is no evidence that this species is using the area for breeding or that it has a particular affinity with the area.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1
Birds	Alca torda	Razorbill	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Although there are a moderate number of records of the manx sheerwater in the AOI there is no evidence that this species is using the area for breeding or that it has a particular affinity with the area.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1

Species Name / Group			Assessment					
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority		
Birds	Fratercula arctica	Puffin	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Very few records of this species (1) in the AOI which migrates through the area in spring and autumn. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2		
Birds	Uria aalge	Common Guillemot	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	This species is known to nest in the AOI and has been recorded in high numbers (100) and will therefore be assessed at the species level.	COMMON GUILLEMOT & PURSUIT & PLUNGE DIVING BIRDS	Tier 1		
Birds	Larus argentatus	Herring Gull	Bonn Convention, Birds Directive, IUCN Red List, English NERC List, UK Cons. Status (Birds)	This species is known to breed and over-winter in the AOI and despite only being recorded in low numbers (5) will be assessed at the species level due to its red conservation status.	HERRING GULL & SURFACE FEEDING BIRDS	Tier 1		
Birds	Larus fuscus	Lesser Black- backed Gull	Bonn Convention, Birds Directive, OSPAR (2008), IUCN Red List, UK Cons. Status (Birds)	This species overwinters in the AOI and has been recorded in moderate numbers (20) it will therefore be assessed at the species level	LESSER-BLACK- BACKED GULL & SURFACE FEEDING BIRDS	Tier 1		
Birds	Larus marinus	Great Black- Backed Gull	Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	This species breeds and nests in the AOI and has been recorded in high numbers (74) it will therefore be assessed at the species level	BLACK-BACKED- GULL & SURFACE FEEDING BIRDS	Tier 1		
Birds	Larus canus	Common Gull	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Few records of this species (11) in the AOI which uses this area to over-winter. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 1		
Birds	Larus ridibundus	Black-headed Gull	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Few records of this species (9) in the AOI although there are large breeding colonies nearby. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 1		
Birds	Hydrocoloeus minutus	Little Gull	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Few records of this species (1) in the AOI which uses migrates through the area in spring / autumn. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 2		
Birds	Larus melanocephalus	Mediterranea n Gull	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Few records of this species (1) in the AOI although there are large breeding colonies nearby. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 2		
Birds	Rissa tridactyla	Black-legged Kittiwake	Bern Convention, Bonn Convention, OSPAR (2008), IUCN Red List, UK Cons. Status (Birds)	Although there are no records of this species in the AOI it is known to breed in nearby areas and there is therefore potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2		
Birds	Stercorarius parasiticus	Arctic Skua	Bern Convention, IUCN Red List, UK Cons. Status (Birds)	Few records of this species (4) in the AOI and this species is likely only to pass through the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 1		
Birds	Stercorarius skua	Great Skua	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Few records of this species (6) in the AOI and this species is likely only to pass through the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	SURFACE FEEDING BIRDS	Tier 1		

Species Name / Group			Assessment				
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority	
Birds	Chlidonias niger	Black Tern	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Few records of this species (1) in the AOI which migrates through the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2	
Birds	Sterna hirundo	Common Tern	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Although there are only a moderate number of records of the common tern in the AOI there is evidence that this species is using the area for breeding . This species is therefore included in the assessment at the species level.	COMMON TERN & PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Sterna paradisaea	Arctic Tern	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Few records of this species (3) in the AOI which migrates through the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Sterna sandvicensis	Sandwich Tern	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Although there are only a small number of this species within the AOI it is known to breed within the AOI and as such will be included in the sensitivity assessment at the species level.	SANDWICH TERN & PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Sterna albifrons	Little Tern	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Although there are no spatial records of this species within the AOI it is known to breed within the AOI and will therefore be included in the sensitivity assessment at the species level.	LITTLE TERN & PURSUIT & PLUNGE DIVING BIRDS	Tier 2	
Birds	Actitis hypoleucos	Common Sandpiper	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Calidris maritima	Purple Sandpiper	Bern Convention, Bonn Convention, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Haematopus ostralegus	Oystercatcher	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Limosa limosa	Black-tailed Godwit	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Morus bassanus	Gannet	Bern Convention, Bonn Convention, IUCN Red List, English NERC List, UK Cons. Status (Birds)	Although there are a high number of records of this species within the AOI there is no evidence of breeding or nesting in this area and the AOI is not thought to have any special significance for this species. Nevertheless is could be impacted by inshore fishing activities.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Phalacrocorax aristotelis	European Shag	Bern Convention, IUCN Red List, UK Cons. Status (Birds)	Although there are no spatial records of this species within the AOI it is known to breed nearby and nest within the AOI and will therefore be included in the assessment at the species level.	EUROPEAN SHAG & PURSUIT & PLUNGE DIVING BIRDS	Tier 2	

Species Name / Group			Assessment				
Species Group	Latin Name	Common Name	Conservation Status	Relevance	Project Model Assessment Unit	Priority	
Birds	Phalacrocorax carbo	Great Cormorant	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	There are a moderate number of records of this species from within the AOI and I it is known to breed and nest within the AOI and will therefore be included in the assessment at the species level.	CORMORANT & PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Anas crecca	Teal	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Anas penelope	Wigeon	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Tadorna tadorna	Shelduck	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Melanitta nigra	Common Scoter	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, English NERC List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Although there are a relatively small number of records of this species within the AOI (11) it is known to overwinter in the AOI and has a high conservation status (red) therefore it will be included in the sensitivity assessment at the species level.	COMMON SCOTER & PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Branta bernicla	Brent Goose	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, English NERC List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Cygnus olor	Mute Swan	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Podiceps cristatus	Great Crested Grebe	Bern Convention, Bonn Convention, IUCN Red List, UK Cons. Status (Birds)	Only a single record of this species in the AOI Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 2	
Birds	Egretta garzetta	Little Egret	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, UK Cons. Status (Birds)	Intertidal species which will not interact with inshore fisheries	Excluded	N/A	
Birds	Gavia immer	Great Northern Diver	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Few records of this species (9) in the AOI which over winters in the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1	
Birds	Gavia stellata	Red Throated Diver	Bern Convention, Bonn Convention, Birds Directive, IUCN Red List, Wildlife & Countryside Act 1981, UK Cons. Status (Birds)	Few records of this species (3) in the AOI which over winters h the area. Nevertheless there is potential for this species to be impacted by the inshore fishing activities in the AOI.	PURSUIT & PLUNGE DIVING BIRDS	Tier 1	

Selected species of conservation importance

Whilst Appendix D shows the final species taken forward in the assessment from a vintage of up to 20 years in age, **Figure E1** shows these same species with Latin names shown; and **Figure E2** shows the same species but for all years data was made available. Only data 1994 onwards was used in the assessment, as shown in the main report.




Figure E1: Location of benthic species selected for risk assessment (Latin names)




Figure E2: Location of selected benthic species from all vintages